

BULLETIN

OF THE

AUSTRALIAN MATHEMATICAL SOCIETY

[\(BibTeX\)](#)

[\(XML metadata\)](#)

[\(RSS feed\)](#)

pp. 87–107: Edward Neuman and József Sándor
On the Ky Fan inequality and related inequalities II.

Volume 72, Number 1
August, 2005

[\(MathReviews\)](#)

[\(Zentralblatt\)](#)

The Australian Mathematical Publishing Assoc. Inc.
Australian National University, ACT 0200, Australia

Printed in Australia by Pirion Printing

Print Post approved - PP229219/00095

ISSN 0004-9727

THE AUSTRALIAN MATHEMATICAL SOCIETY

President: M.G. Cowling
Department of Pure Mathematics,
The University of New South Wales,
Sydney NSW 2052, Australia.

Secretary: E.J. Billington
Department of Mathematics,
The University of Queensland,
Queensland 4072, Australia.

Treasurer: A. Howe
Department of Mathematics,
The Australian National University,
Canberra ACT 0200, Australia

Membership and correspondence.

Applications for membership, notices of change of address or title or position, members' subscriptions and correspondence related to accounts should be sent to the Treasurer. Correspondence about the distribution of the Society's BULLETIN, GAZETTE, and JOURNALS, and orders for back numbers should be sent to the Treasurer. All other correspondence should be sent to the Secretary.

The Bulletin.

The Bulletin of the Australian Mathematical Society began publication in 1969. Normally two volumes of three numbers are published annually. The BULLETIN is published for the Australian Mathematical Society by the Australian Mathematical Publishing Association Inc.

Editor: Alan S. Jones
Deputy Editor: Graeme A. Chandler
Department of Mathematics,
The University of Queensland,
Queensland 4072, Australia.

ASSOCIATE EDITORS

Robert S. Anderssen	B.D. Craven	B.D. Jones	M. Murray
R. Bartnik	Brian A. Davey	Owen D. Jones	J.H. Rubinstein
Elizabeth J. Billington	J.R. Giles	G.I. Lehrer	Jamie Simpson
G. Cairns	J.A. Hempel	K.L. McAvaney	Brailey Sims
J. Clark	B.D. Hughes	A.G.R. McIntosh	Ross Street
G.L. Cohen	G. Ivanov	Terry Mills	R.P. Sullivan
John Cossey	B.R.F. Jefferies	Sidney A. Morris	H.B. Thompson
N.S. Trudinger		A.J. van der Poorten	

©Copyright Statement Where necessary, permission to photocopy for internal or personal use or the internal or personal use of specific clients is granted by the Treasurer, Australian Mathematical Publishing Association, Inc., for libraries and other users registered with the Copyright Clearance Center (CCC), provided that the base fee of \$A2.00 per copy of article is paid directly to CCC, 21 Congress Street, Salem, MA 01970, U.S.A. Special requests should be addressed to the Treasurer, Australian Mathematical Publishing Association, Inc., School of Mathematical Sciences, ANU, Canberra ACT 0200 Australia. Serial-fee code: 0004-9727/05 \$A2.00 + 0.00.

INFORMATION FOR AUTHORS

The Bulletin of the Australian Mathematical Society aims at quick publication of original research in all branches of mathematics. The Editors receive more than three times as much material as can be published in the BULLETIN; many meritorious papers can, therefore, not be accepted. Authors are asked to avoid, as far as possible the use of mathematical symbols in the title. Manuscripts are accepted for review with the understanding that the same work is not concurrently submitted elsewhere.

To ensure speedy publication, editorial decisions on acceptance or otherwise are taken quickly, normally within a month of receipt of the paper. Papers are accepted only after a careful evaluation by the Editor and an Associate Editor or other expert in the field. As even minor revisions are generally not permitted, authors should read carefully all the details listed below. For a paper to be acceptable for publication, not only should it contain new and interesting results but also

- (i) the exposition should be clear and attractive;
- (ii) the manuscript should be in publishable form, without revision.

Authors should submit three clean, high quality copies to

The Editorial Office, Bulletin of the Australian Mathematical Society,
Department of Mathematics, The University of Queensland,
Queensland 4072, Australia.

Unless requested at the time, material submitted to the BULLETIN will usually not be returned.

EDITORIAL POLICY

1. References. Arrange references alphabetically (by surname of the first author) and cite them numerically in the text. Ensure the accuracy of the references: authors' names should appear as in the work quoted. Include in the list of references only those works cited, and avoid citing works which are "in preparation" or "submitted". Where the work cited is not readily accessible (for example, a preprint) a photocopy of the title page and relevant sections of the copy that you have used should be included with your submission.

2. Abstracts.

1. Each paper must include an abstract of not more than 200 words, which should contain a brief but informative summary of the contents of the paper, but no inessential details.
2. The abstract should be self-contained, but may refer to the title.
3. Specific references (by number) to a section, proposition, equation or bibliographical item should be avoided.

3. Subject Classification. Authors should include in their papers one or more classification numbers, following the 2000 Mathematics Subject Classification. Details of this scheme can be found in each Annual Index of Mathematical Reviews or on the web at <http://www.ams.org/msc> .

4. Abstracts of Ph.D. Theses. The Bulletin endeavours to publish abstracts of all accepted Australasian Ph.D. theses in mathematics. One restriction, however, is that the abstract must be received by the Editor within 6 months of the degree being approved.

5. Electronic Manuscripts. The Bulletin is produced using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-T}\mathcal{E}\mathcal{X}$. Authors who are able to do so are invited to prepare their manuscripts using $\text{T}\mathcal{E}\mathcal{X}$. (We accept Plain $\text{T}\mathcal{E}\mathcal{X}$, $\mathcal{A}\mathcal{M}\mathcal{S}\text{-T}\mathcal{E}\mathcal{X}$ or $\text{L}\mathcal{A}\text{T}\mathcal{E}\mathcal{X}$.) Hard copy only should be submitted for assessment, but if the paper is accepted the author will be asked to send the text on an IBM PC compatible diskette or via e-mail to ams@maths.uq.edu.au. [Typed manuscripts are, of course, still acceptable.]

Bulletin of the Australian Mathematical Society

A note on the lattice of density preserving maps	
Sejal Shah and T.K. Das	1
A strong excision theorem for generalised Tate cohomology	
N. Mramor Kosta	7
Linear geometries on the Moebius strip: a theorem of Skornyakov type	
Rainer Löwen and Burkhard Polster	17
Div-curl type theorems on Lipschitz domains	
Zengjian Lou	31
A nonlinear map for midpoint locally uniformly rotund renorming	
S. Lajara and A.J. Pallarés	39
A remarkable continued fraction	
David Angell and Michael D. Hirschhorn	45
A new variational method for the $p(x)$-Laplacian equation	
Marek Galewski	53
Boundary unique continuation theorems under zero Neumann boundary conditions	
Xiangxing Tao and Songyan Zhang	67
On the Ky Fan inequality and related inequalities II	
Edward Neuman and József Sándor	87
Finite presentability of some metabelian Hopf algebras	
Dessislava H. Kochloukova	109
On the monotonicity properties of additive representation functions	
Yong-Gao Chen, András Sárközy, Vera T. Sós and MinTang	129
Generation of diagonal acts of some semigroups of transformations and relations	
Peter Gallagher and Nik Ruškuc	139
Subalgebras of free restricted Lie algebras	
R.M. Bryant, L.G. Kovács and Ralph Stöhr	147
A multiple character sum evaluation	
Dae San Kim	157
Implicit vector equilibrium problems via nonlinear scalarisation	
Jun Li and Nan-Jing Huang	161

ABSTRACTS OF AUSTRALASIAN Ph.D. THESES

Numerical methods for quantitative finance	
Jamie Alcock	173

ON THE KY FAN INEQUALITY AND RELATED INEQUALITIES II

EDWARD NEUMAN AND JÓZSEF SÁNDOR

Ky Fan type inequalities for means of two or more variables are obtained. Refinements and improvements of known inequalities are derived. Applications to symmetric elliptic integrals of the first and second kind are also included.

1. INTRODUCTION AND NOTATION

This paper deals with the Ky Fan type inequalities for means of two and more variables and is a continuation of our earlier work [15].

Let $X = (x_1, \dots, x_n)$ be an n -tuple of positive numbers. In what follows the symbols A_n , G_n , and H_n will stand for the unweighted arithmetic, geometric, and harmonic means of X , respectively, that is,

$$A_n = \frac{1}{n} \sum_{i=1}^n x_i, \quad G_n = \left(\prod_{i=1}^n x_i \right)^{1/n}, \quad H_n = \frac{n}{\sum_{i=1}^n (1/x_i)}.$$

We shall always assume that $x_i \leq 1/2$ for $i = 1, 2, \dots, n$. Also, let $X' = 1 - X = (1 - x_1, \dots, 1 - x_n)$. Throughout the sequel the unweighted arithmetic, geometric, and harmonic means of X' will be denoted by A'_n , G'_n , and H'_n , respectively.

The classical Ky Fan inequality reads as follows

$$(1.1) \quad \frac{G_n}{G'_n} \leq \frac{A_n}{A'_n}$$

(see, for example, [3]). A companion inequality to (1.1) has been obtained by Wang and Wang in [24], and it states that

$$(1.2) \quad \frac{H_n}{H'_n} \leq \frac{G_n}{G'_n}.$$

These two inequalities have stimulated the interest of several researchers, which resulted in many new results, refinements and improvements. The interested reader is referred to [1, 2, 5, 6, 7, 9, 11, 15, 16, 17, 20, 21, 22].

Received 1st March, 2005

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/05 \$A2.00+0.00.

In the following by $w = (w_1, \dots, w_n)$, $w_i \geq 0$ ($1 \leq i \leq n$), $w_1 + \dots + w_n = 1$ denotes the n -tuple of weights. The weighted arithmetic, geometric, and harmonic means of $X \in \mathbb{R}_+^n$ with the weights w , denoted by $A(X, w)$, $G(X, w)$ and $H(X, w)$, are defined in the usual way:

$$(1.3) \quad \begin{aligned} A(X, w) &= \sum_{i=1}^n w_i x_i, & G(X, w) &= \prod_{i=1}^n x_i^{w_i}, \\ H(X, w) &= \frac{1}{\sum_{i=1}^n w_i (1/x_i)}. \end{aligned}$$

Other weighted means used in this paper include the logarithmic mean $L(X, w)$ and the identric mean $I(X, w)$. Both means admit integral representations which are given below. Let

$$E_{n-1} = \{(u_1, \dots, u_{n-1}) : u_i \geq 0 \ (1 \leq i \leq n-1), \ u_1 + \dots + u_{n-1} \leq 1\}$$

denote the Euclidean simplex, and for $(u_1, \dots, u_{n-1}) \in E_{n-1}$, put

$$u_n := 1 - (u_1 + \dots + u_{n-1}).$$

Throughout the paper $n \geq 2$ and μ is an arbitrary probability measure on E_{n-1} . The natural weights w_i ($1 \leq i \leq n$) of the measure μ are defined by

$$(1.4) \quad w_i = \int_{E_{n-1}} u_i d\mu(u),$$

where $d\mu(u) = \mu(u) du_1 \cdots du_{n-1}$ and $u = (u_1, \dots, u_n)$ (see, for example, [4]). The weighted logarithmic mean of X is defined as

$$(1.5) \quad L(X, w) = \left[\int_{E_{n-1}} (u \cdot X)^{-1} d\mu(u) \right]^{-1},$$

where $u \cdot X = u_1 x_1 + \dots + u_n x_n$ is the dot product of u and X (see, for example, [18]). The weighted identric mean is given by

$$(1.6) \quad I(X, w) = \exp \left[\int_{E_{n-1}} \ln(u \cdot x) d\mu(u) \right]$$

(see [18]).

An important probability measure on E_{n-1} is the Dirichlet measure $\mu_b(u)$, where $b \in \mathbb{R}_+^n$. It is defined by

$$(1.7) \quad \mu_b(u) = \frac{1}{B(b)} \prod_{i=1}^n u_i^{b_i-1},$$

where $B(\cdot)$ stands for the multivariate beta function (see [4]). It is known that the natural weights w_i of μ_b are given by

$$(1.8) \quad w_i = b_i/c$$

($1 \leq i \leq n$) where $c = b_1 + \cdots + b_n$ (see [4]).

The weighted geometric mean also admits an integral representation

$$(1.9) \quad G(X, w) = \left[\int_{E_{n-1}} (u \cdot X)^{-c} d\mu_b(u) \right]^{-1/c}.$$

This follows from [4, (6.6–6)]. It is known that the weighted means mentioned above satisfy the inequalities

$$(1.10) \quad \begin{aligned} H(X, w) &\leq G(X, w) \leq L(X, w) \leq I(X, w) \\ &\leq A(X, w) \leq M_p(X, w) \end{aligned}$$

($p \geq 1$) where

$$(1.11) \quad M_p(X, w) = \begin{cases} \left(\sum_{i=1}^n w_i x_i^p \right)^{1/p}, & p \neq 0 \\ G(X, w), & p = 0 \end{cases}$$

is the weighted power mean of order p of $X \in \mathbb{R}_+^n$. Some of the means which appear in (1.10) satisfy the Ky Fan type inequalities. For instance, Sándor and Trif [22] have proven that

$$(1.12) \quad \frac{G(X, w)}{G'(X, w)} \leq \frac{I(X, w)}{I'(X, w)} \leq \frac{A(X, w)}{A'(X, w)}.$$

(See also [21] for a special case of (1.12)).

Let the letters G , L , and A stand for the unweighted geometric, logarithmic, and arithmetic means, respectively, of two positive numbers x and y . Recall that

$$L \equiv L(x, y) = \begin{cases} \frac{x-y}{\ln(x/y)}, & x \neq y \\ x, & x = y. \end{cases}$$

In [16] the authors have proven that

$$(1.13) \quad \frac{G}{G'} \leq \frac{L}{L'} \leq \frac{A}{A'}$$

($x, y \leq 1/2$). In addition to these inequalities, results involving differences of the reciprocals of means have been obtained. The following result

$$(1.14) \quad \begin{aligned} \frac{1}{H'(X, w)} - \frac{1}{H(X, w)} &\leq \frac{1}{L'(X, w)} - \frac{1}{L(X, w)} \\ &\leq \frac{1}{A'(X, w)} - \frac{1}{A(X, w)} \end{aligned}$$

is established in [15]. The middle term in (1.14) provides a refinement of an earlier result of Sándor [20]. Alzer [2] has shown that

$$(1.15) \quad \frac{1}{H'(X, w)} - \frac{1}{H(X, w)} \leq \frac{1}{G'(X, w)} - \frac{1}{G(X, w)}.$$

The same author (see [1]) also proved that

$$(1.16) \quad \frac{1}{H'_n} - \frac{1}{H_n} \leq \frac{1}{G'_n} - \frac{1}{G_n} \leq \frac{1}{A'_n} - \frac{1}{A_n}.$$

The goal of this paper is to obtain Ky Fan type inequalities for the means mentioned in this section and also for other means discussed in Section 5. In Section 2, we give some lemmas which will be used throughout the sequel. Therein we introduce the so-called Ky Fan rules which, among other things, allow us to obtain inequalities for the differences of reciprocals of means. For instance, an application of Lemma 2.1, part (i) to (1.1) and (1.2) gives immediately Alzer's result (1.16). New Ky Fan type inequalities for the multivariate means are obtained in Section 3. Two refinements of the inequality (1.1) are presented in Section 4. The next section is devoted to the study of the Ky Fan type inequalities for certain means derived from A_n , G_n and H_n . Section 6, deals with the Ky Fan type inequalities for the unweighted means of two variables.

2. LEMMAS

The lemmas presented in this section will be used in the remaining parts of this paper. We shall refer to the inequalities (2.1)–(2.3) of Lemma 2.1 as the Ky Fan rules. Our first result reads as follows.

LEMMA 2.1. *Let a, b, a' and b' be positive numbers.*

(i) *If $a \leq b$ and $a/a' \leq b/b' \leq 1$ or if $b \leq a$ and $1 \leq a/a' \leq b/b'$, then*

$$(2.1) \quad \frac{1}{a'} - \frac{1}{a} \leq \frac{1}{b'} - \frac{1}{b}.$$

(ii) *If $a' \leq b'$ and $a/a' \leq b/b'$, then*

$$(2.2) \quad aa' \leq bb'.$$

(iii) *If $a/a' \leq b/b'$, then*

$$(2.3) \quad \frac{a}{a'} \leq \frac{a+b}{a'+b'} \leq \frac{b}{b'}$$

and

$$(2.4) \quad \frac{a}{a'} \leq \frac{\sqrt{ab}}{\sqrt{a'b'}} \leq \frac{b}{b'}.$$

PROOF: For the proof of (2.1) consider the case when $a \leq b$ and $a/a' \leq b/b' \leq 1$. It follows that $1/b \leq 1/a$ and also that $1 - b/b' \leq 1 - a/a'$. Multiplying corresponding sides of the last two inequalities gives the desired result. The remaining case can be established in the same way. For the proof of (2.2) we use the assumptions to obtain $a'/b' \leq b'/a'$ and $b'/a' \leq b/a$. Combining the last two inequalities we obtain $a'/b' \leq b/a$. Hence (2.2) follows. We shall prove now the inequalities (2.3). We have $ab' \leq a'b$. Adding aa' to both sides we obtain $a(a'+b') \leq a'(a+b)$. Hence the first inequality in (2.3) follows. The second one is obtained from $ab' \leq a'b$ after adding to both sides bb' . The first inequality in (2.4) is derived from $a/a' \leq b/b'$. Multiplying both sides by a/a' and next extracting square roots of the resulting expressions gives the assertion. The second inequality in (2.4) is obtained from $a/a' \leq b/b'$ by multiplying both sides by b/b' and followed by the root extraction. \square

LEMMA 2.2. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be a convex (concave) function. Then the function $g : [0, 1/2] \rightarrow \mathbb{R}$, given by $g(x) = f(x) + f(1-x)$ is a decreasing (increasing) function on its domain.*

PROOF: Assume that the numbers x and y are such that $0 < x < y \leq 1/2$. Then $x < y \leq 1 - y < 1 - x$, that is, both numbers y and $1 - y$ can be written as a convex combination of x and $1 - x$, that is, $y = \alpha x + \beta(1 - x)$ and $1 - y = \beta x + \alpha(1 - x)$ ($\alpha, \beta \geq 0$, $\alpha + \beta = 1$). Making use of the definition of the function g together with the assumption of convexity of f we obtain

$$\begin{aligned} g(y) &= f(y) + f(1 - y) \leq \alpha f(x) + \beta f(1 - x) + \beta f(x) + \alpha f(1 - x) \\ &= f(x) + f(1 - x) = g(x). \end{aligned}$$

This completes the proof. \square

LEMMA 2.3. *Let $0 < x < y \leq 1/2$ and let $x' = 1 - x$ and $y' = 1 - y$. If $p = c \ln(x/y)$ and $q = c \ln(x'/y')$, where $c > 0$, then the following inequalities*

$$(2.5) \quad \frac{\sinh q}{q} < \frac{\sinh p}{p},$$

$$(2.6) \quad \cosh q < \cosh p,$$

and

$$(2.7) \quad \frac{\tanh p}{p} < \frac{\tanh q}{q}$$

are valid.

PROOF: We shall show first that the numbers p and q satisfy the inequalities

$$(2.8) \quad -p > q > 0.$$

We have $0 < x < y \leq y' < x' < 1$. This in turn implies that the logarithmic means of (x, y) and (x', y') satisfy the inequality $L(x, y) < L(x', y')$. Thus

$$\frac{y-x}{\ln(y/x)} < \frac{x'-y'}{\ln(x'/y')} = \frac{y-x}{\ln(x'/y')}.$$

Since $y-x > 0$, $\ln(y/x) > \ln(x'/y')$. Multiplying both sides by $c > 0$ we obtain (2.8). To complete the proof of inequalities (2.5)–(2.7) we utilise the fact that the functions $\sinh t/t$, $\cosh t$ and $\tanh t/t$ are even functions on \mathbb{R} and also that the first two are strictly increasing on \mathbb{R}_+ while the third one is strictly decreasing on the stated domain. Thus we have

$$\frac{\sinh(-p)}{-p} = \frac{\sinh p}{p} > \frac{\sinh q}{q},$$

$\cosh(-p) = \cosh p > \cosh q$ and

$$\frac{\tanh(-p)}{-p} = \frac{\tanh p}{p} < \frac{\tanh q}{q}.$$

The proof is complete. □

We shall also need the following.

LEMMA 2.4. *Let f be a continuous function on the interval $[a, b]$ and assume that all derivatives $f^{(\ell)}(m)$ ($\ell = 1, 2, \dots$) exist at $m = (a+b)/2$. Then*

$$(2.9) \quad \frac{1}{b-a} \int_a^b f(t) dt = \frac{1}{2} [f(a) + f(b)] - \sum_{k=1}^{\infty} \frac{2k}{(2k+1)!} \left(\frac{b-a}{2}\right)^{2k} f^{(2k)}(m).$$

PROOF: Integrating Taylor's expansion for the function f about the point $t = m$

$$(2.10) \quad f(t) = f(m) + \sum_{k=1}^{\infty} \frac{(t-m)^k}{k!} f^{(k)}(m)$$

we obtain

$$(2.11) \quad \frac{1}{b-a} \int_a^b f(t) dt = f(m) + \sum_{k=1}^{\infty} \frac{1}{(2k+1)!} \left(\frac{b-a}{2}\right)^{2k} f^{(2k)}(m).$$

Next we substitute $t = a$ and $t = b$ into (2.10) and add up the resulting expressions to obtain

$$(2.12) \quad \frac{1}{2} [f(a) + f(b)] = f(m) + \sum_{k=1}^{\infty} \frac{1}{(2k)!} \left(\frac{b-a}{2}\right)^{2k} f^{(2k)}(m).$$

Eliminating $f(m)$ between (2.11) and (2.12) gives the desired result (2.9). □

3. INEQUALITIES INVOLVING MEANS OF SEVERAL VARIABLES

In this section we give several Ky Fan type inequalities for means of several variables. For the sake of notation we shall write A for the weighted arithmetic mean $A(X, w)$, L for the weighted logarithmic mean $L(X, w)$, et cetera. Recall that $X = (x_1, \dots, x_n) \in \mathbb{R}_+^n$ where $x_i \leq 1/2$, $1 \leq i \leq n$.

Our first result reads as follows.

THEOREM 3.1. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be a convex function. Then*

$$(3.1) \quad \begin{aligned} f(H) + f(1 - H) &\geq f(G) + f(1 - G) \geq f(L) + f(1 - L) \\ &\geq f(I) + f(1 - I) \geq f(A) + f(A'). \end{aligned}$$

Inequalities (3.1) are reversed if f is a concave function on $[0, 1]$. If $F : (0, 1] \rightarrow \mathbb{R}_+$ is a log-convex function, then

$$(3.2) \quad \begin{aligned} F(H)F(1 - H) &\geq F(G)F(1 - G) \geq F(L)F(1 - L) \\ &\geq F(I)F(1 - I) \geq F(A)F(A'). \end{aligned}$$

Inequalities (3.2) are reversed if F is a log-concave function and they become equalities if $F(x) = (1 - x)/x$.

PROOF: For the proof of the inequalities (3.1) we apply Lemma 2.2 to the five first members of (1.1) together with an obvious identity $A + A' = 1$. Inequalities (3.2) follow from (3.1) by letting $f(x) = \ln F(x)$. The last statement of the theorem is obvious. \square

COROLLARY 3.2. *The following inequalities*

$$(3.3) \quad \frac{1}{H} + \frac{1}{1 - H} \geq \frac{1}{G} + \frac{1}{1 - G} \geq \frac{1}{L} + \frac{1}{1 - L} \geq \frac{1}{I} + \frac{1}{1 - I} \geq \frac{1}{A} + \frac{1}{A'},$$

$$(3.4) \quad \frac{1}{G} + \frac{1}{G'} \geq \frac{1}{L} + \frac{1}{1 - L} \geq \frac{1}{I} + \frac{1}{1 - I} \geq \frac{1}{A} + \frac{1}{A'},$$

$$(3.5) \quad \frac{1}{H} + \frac{1}{H'} \geq \frac{1}{G} + \frac{1}{1 - G} \geq \frac{1}{L} + \frac{1}{1 - L} \geq \frac{1}{I} + \frac{1}{1 - I} \geq \frac{1}{A} + \frac{1}{A'},$$

are valid.

PROOF: Inequalities (3.3) follow immediately from (3.1) by letting $f(x) = 1/x$. Since $G + G' \leq A + A' = 1$, $G' \leq 1 - G$ and $1/G + 1/(1 - G) \leq 1/G + 1/G'$. This in conjunction with (3.3) gives the chain of inequalities (3.4). Using the same argument as above, we have $H' \leq 1 - H$. Hence $1/H + 1/H' \geq 1/H + 1/(1 - H)$. Combining this with (3.3) yields (3.5) \square

A multiplicative version (with refinements) of the Ky Fan inequality is contained in the following.

COROLLARY 3.3. *We have*

$$(3.6) \quad H(1 - H) \leq G(1 - G) \leq L(1 - L) \leq I(1 - I) \leq AA',$$

$$(3.7) \quad GG' \leq L(1 - L) \leq I(1 - I) \leq AA',$$

and

$$(3.8) \quad \begin{aligned} \Gamma(H)\Gamma(1 - H) &\geq \Gamma(G)\Gamma(1 - G) \geq \Gamma(L)\Gamma(1 - L) \\ &\geq \Gamma(I)\Gamma(1 - I) \geq \Gamma(A)\Gamma(A'), \end{aligned}$$

where Γ stands for Euler's gamma function.

PROOF: Inequalities (3.6) follow from (3.1) by letting $f(t) = \ln t$. Taking into account that $G' \leq 1 - G$ we obtain (3.7) from (3.6). It is well-known that the gamma function $\Gamma(x)$ is log-convex for $x > 0$ (see, for example, [4]). Letting $F = \Gamma$ in (3.2) gives the inequalities (3.8). \square

In [19], Sándor has proven that the function $f(x) = [\Gamma(x)]^{1/x}$ is convex for $x > 0$. This in conjunction with (3.1) gives more Ky Fan type inequalities. We omit further details.

Before we shall state the next result let us introduce more notation. Also, we shall use the concept of an integral average of a function. For $Y = (y_1, \dots, y_n) \in \mathbb{R}^n$, let $C = [\min(Y), \max(Y)]$ with $\min(Y) < \max(Y)$. Further, let $f : C \rightarrow \mathbb{R}$ be a continuous function and let μ be a probability measure on the Euclidean simplex E_{n-1} . We define

$$F(\mu; Y) = \int_{E_{n-1}} f(u \cdot Y) d\mu(u).$$

Application of the mean-value theorem for integrals gives $F(\mu; Y) = f(\xi \cdot Y)$, where $\xi = (\xi_1, \dots, \xi_{n-1}, 1 - \xi_1 - \dots - \xi_{n-1})$ with $(\xi_1, \dots, \xi_{n-1}) \in E_{n-1}$. Thus $F(\mu; Y)$ can be regarded as the average value of the function f . When $\mu = \mu_b$, the average $F(\mu_b, Y) \equiv F(b; Y)$ is studied in detail in Carlson's monograph [4] and is called the Dirichlet average of the function f .

Also, recall that a function f is said to be n -convex on C if $f[z_0, \dots, z_n] \geq 0$ for all choices of $n+1$ points z_0, \dots, z_n in C . Here $f[z_0, \dots, z_n]$ stands for the n -th order divided difference of f .

The following refinement of Levinson's inequality (see [11]) appears in [14, Theorem 3.4]. Let $f : (0, 1] \rightarrow \mathbb{R}$ be a 3-convex function. Then

$$(3.9) \quad f(A') - f(A) \leq F(\mu; X') - F(\mu; X) \leq \sum_{i=1}^n w_i [f(x'_i) - f(x_i)].$$

Inequalities (3.9) are reversed if f is a 3-concave function on $(0, 1]$.

The integral average of the power function $f(t) = t^p$ ($t > 0, p \in \mathbb{R}$) will be denoted by $R_p(\mu; \cdot)$ and when $\mu = \mu_b$, the Dirichlet measure on E_{n-1} , we shall simply write $R_p(b; \cdot)$ instead of $R_p(\mu_b; \cdot)$. The latter average is also called the R -hypergeometric function. Its importance in the theory of special functions is well documented in Carlson's monograph [4]. If $\alpha > 0$ and if $\alpha' := b_1 + \cdots + b_n - \alpha > 0$, then the R -hypergeometric function $R_{-\alpha}$ admits another integral representation (see [4, (6.8–6)])

$$R_{-\alpha}(b; X) = \frac{1}{B(\alpha, \alpha')} \int_0^\infty t^{\alpha'-1} \prod_{i=1}^n (t + x_i)^{-b_i} dt.$$

We shall use inequalities (3.9) to obtain bounds for the symmetric elliptic integrals of the first and the second kind. They are denoted by R_F and R_G , respectively, and defined as follows

$$R_F(x, y, z) = \frac{1}{2} \int_0^\infty [(t+x)(t+y)(t+z)]^{-1/2} dt$$

and

$$R_G(x, y, z) = \frac{1}{4} \int_0^\infty [(t+x)(t+y)(t+z)]^{-1/2} \left(\frac{x}{t+x} + \frac{y}{t+y} + \frac{z}{t+z} \right) t dt$$

(see [4]). We assume that x, y, z are positive numbers and at most one of them is 0.

We have the following.

PROPOSITION 3.4. *Let μ be an arbitrary probability measure on E_{n-1} and let $X \in \mathbb{R}_+^n$ with $x_i \leq 1/2$ for $1 \leq i \leq n$. If $0 < p < 1$ or if $p > 2$, then*

$$(3.10) \quad (A')^p - A^p < R_p(\mu; X') - R_p(\mu; X) < (M'_p)^p - M_p^p.$$

Inequalities in (3.10) are reversed if either $p < 0$ or if $1 < p < 2$ and they become equalities if $p = 0$ or $p = 1$ or $p = 2$.

PROOF: Let $f(t) = t^p$ ($t > 0, p \in \mathbb{R}$). It is easy to see that f is 3-convex if $0 < p < 1$ or if $p > 2$ and f is 3-concave if either $p < 0$ or $1 < p < 2$. The inequalities (3.10) follow from (3.9). If $p = 0$, then all three members in (3.10) are equal to zero and they are all equal to $A' - A$ if either $p = 1$ or $p = 2$. \square

COROLLARY 3.5. *The symmetric elliptic integrals R_F and R_G satisfy the Ky Fan type inequalities*

$$(3.11) \quad (A')^{-1/2} - A^{-1/2} > R_F(X') - R_F(X) > (M'_{-1/2})^{-1/2} - M_{-1/2}^{-1/2}$$

and

$$(3.12) \quad (A')^{1/2} - A^{1/2} < R_G(X') - R_G(X) < (M'_{1/2})^{1/2} - M_{1/2}^{1/2},$$

where $X = (x_1, x_2, x_3)$ with $0 < x_i \leq 1/2, 1 \leq i \leq 3$.

PROOF: We use Proposition 3.4 with $\mu = \mu_b$ where $b = (1/2, 1/2, 1/2)$ and $p = -1/2$ or $p = 1/2$, respectively, because $R_F(\cdot) = R_{-1/2}(b; \cdot)$ and $R_G(\cdot) = R_{1/2}(b; \cdot)$ (see [4]). \square

For more bounds for the symmetric elliptic integrals the interested reader is referred to [13].

We close this section with two Ky Fan type inequalities for the differences of reciprocals of certain multivariate means.

The following inequality

$$\frac{L}{L'} \leq \frac{I}{I'}$$

is established in [7]. This result was known to the authors of this paper prior to the publication of [7]. Application of Lemma 2.1, part (i) to the last inequality and the second inequality in (1.12) gives

$$(3.13) \quad \frac{1}{L'} - \frac{1}{L} \leq \frac{1}{I'} - \frac{1}{I} \leq \frac{1}{A'} - \frac{1}{A}.$$

A refinement of the first inequality in (1.16) is provided by

$$(3.14) \quad \frac{1}{H'_n} - \frac{1}{H_n} \leq \frac{1}{J'_n} - \frac{1}{J_n} \leq \frac{1}{G'_n} - \frac{1}{G_n},$$

where

$$J_n := \frac{I_n(Y)}{G_n(Y)} H_n$$

with $Y = (y_1, \dots, y_n)$, $y_i = x_1 \cdot \dots \cdot x_{i-1} x_{i+1} \cdot \dots \cdot x_n$ ($1 \leq i \leq n$). In [15] the authors have proven that $H_n \leq J_n \leq G_n$ and also that

$$(3.15) \quad \frac{H_n}{H'_n} \leq \frac{J_n}{J'_n} \leq \frac{G_n}{G'_n}.$$

Application of Lemma 2.1, part (i) to (3.15) gives immediately the inequalities (3.14).

4. TWO REFINEMENTS OF THE KY FAN INEQUALITY

The goal of this section is to establish two refinements of the Ky Fan inequality (1.1).

Let C be an interval in \mathbb{R} . Recall that a function $f : C \rightarrow \mathbb{R}$ is said to be Jensen's convex (or J -convex) on C if Jensen's inequality

$$f\left(\frac{1}{n} \sum_{i=1}^n y_i\right) \leq \frac{1}{n} \sum_{i=1}^n f(y_i)$$

is satisfied for all y_i ($1 \leq i \leq n$) in C .

We need the following.

PROPOSITION 4.1. Let $x_i \in C$, $1 \leq i \leq n+1$ and let $f : C \rightarrow \mathbb{R}$ be a J -convex function. If $\bar{x}_n = (x_{n+1} + (n-1)A_{n+1})/n$, then

$$(4.1) \quad \begin{aligned} 2f(A_{n+1}) &\leq f(A_n) + f(\bar{x}_n) \\ &\leq \frac{f(x_1) + \cdots + f(x_{n+1}) + (n-1)f(A_{n+1})}{n}. \end{aligned}$$

Also, if $x_n^* = (A_{n+1} + (n-1)x_{n+1})/n$, then

$$(4.2) \quad \begin{aligned} nf(A_{n+1}) &\leq (n-1)f(A_n) + f(x_n^*) \\ &\leq \frac{(n-1)[f(x_1) + \cdots + f(x_{n+1})] + f(A_{n+1})}{n}. \end{aligned}$$

PROOF: For the proof of the inequalities (4.1), let us notice that $A_{n+1} = (A_n + \bar{x}_n)/2$. Using Jensen's inequality three times we obtain

$$\begin{aligned} f(A_{n+1}) &= f\left(\frac{A_n + \bar{x}_n}{2}\right) \leq \frac{f(A_n) + f(\bar{x}_n)}{2} \\ &\leq \frac{1}{2} \left[\frac{f(x_1) + \cdots + f(x_n)}{n} + \frac{f(x_{n+1}) + (n-1)f(A_{n+1})}{n} \right]. \end{aligned}$$

Hence the assertion follows. In the proof of (4.2) we shall utilise the following identity $A_{n+1} = [(n-1)A_n + x_n^*]/n$. Again, we appeal to Jensen's inequality to obtain

$$\begin{aligned} f(A_{n+1}) &= f\left(\frac{(n-1)A_n + x_n^*}{n}\right) \leq \frac{(n-1)f(A_n) + f(x_n^*)}{n} \\ &\leq \frac{1}{n} \left[(n-1) \frac{f(x_1) + \cdots + f(x_n)}{n} + \frac{f(A_{n+1}) + (n-1)f(x_{n+1})}{n} \right]. \end{aligned}$$

The desired result now follows. \square

The main result of this section is contained in the following.

THEOREM 4.2. Let $X \in \mathbb{R}_+^{n+1}$ with $x_i \leq 1/2$ for $1 \leq i \leq n+1$ and let the numbers \bar{x}_n and x_n^* have the same meaning as in Proposition 4.1. Then

$$(4.3) \quad \left(\frac{A'_{n+1}}{A_{n+1}}\right)^2 \leq \frac{A'_n \bar{x}'_n}{A_n \bar{x}_n} \leq \left(\frac{G'_{n+1}}{G_{n+1}}\right)^{1+1/n} \left(\frac{A'_{n+1}}{A_{n+1}}\right)^{1-1/n},$$

where $\bar{x}'_n = 1 - x_n$. Also,

$$(4.4) \quad \frac{A'_{n+1}}{A_{n+1}} \leq \left(\frac{A'_n}{A_n}\right)^{1-1/n} \left(\frac{x'_n}{x_n^*}\right)^{1/n} \leq \left(\frac{G'_{n+1}}{G_{n+1}}\right)^{1-1/n^2} \left(\frac{A'_{n+1}}{A_{n+1}}\right)^{1/n^2}.$$

where $x'_n = 1 - x_n^*$.

REMARK. It is easy to see, using the Ky Fan inequality (1.1) that the third members in (4.3) and (4.4) are bounded above by $(G'_{n+1}/G_{n+1})^2$ and by G'_{n+1}/G_{n+1} , respectively. Thus the inequalities (4.3) and (4.4) provide refinements of (1.1).

PROOF: We shall use Proposition 4.1 with $C = (0, 1/2]$ and $f(x) = \ln(1-x)/x$. Easy computations show that (4.3) follows from (4.1) while (4.4) is a consequence of (4.2). We omit further details. \square

COROLLARY 4.3. *The following inequalities*

$$(4.5) \quad \frac{A'_n}{A_n} \leq \left(\frac{G'_{n+1}}{G_{n+1}} \right)^{1+1/n} \left(\frac{A'_{n+1}}{A_{n+1}} \right)^{1-1/n}$$

and

$$(4.6) \quad \left(\frac{A'_n}{A_n} \right)^{1-1/n} \leq \left(\frac{G'_{n+1}}{G_{n+1}} \right)^{1-1/n^2} \left(\frac{A'_{n+1}}{A_{n+1}} \right)^{1/n^2}$$

are valid.

PROOF: It follows from the definitions of the numbers \bar{x}_n and x_n^* that both are positive and not bigger than $1/2$. This in turn implies that $\bar{x}'_n \geq \bar{x}_n$ and $x_n^{*'} \geq x_n^*$. The inequalities (4.5) and (4.6) now follow from (4.3) and (4.4), respectively. \square

Before we state the last result of this section, let us introduce more notation. We define

$$a_n = \frac{1}{A_n} - \frac{1}{A'_n} \text{ and } h_n = \frac{1}{H_n} - \frac{1}{H'_n}.$$

COROLLARY 4.4. *We have*

$$(4.7) \quad 2a_{n+1} \leq a_n + \tilde{x}_n \leq \frac{n+1}{n}h_{n+1} + \frac{n-1}{n}a_{n+1},$$

where $\tilde{x}_n = 1/\bar{x}_n - 1/\bar{x}'_n$.

PROOF: Inequalities (4.7) follow easily from (4.1) when $C = (0, 1/2]$ and $f(x) = 1/x - 1/(1-x)$. \square

In [20] the author has proven that $a_{n+1} \leq h_{n+1}$ (see also (1.16)). This in conjunction with (4.7) gives

$$a_{n+1} \leq (a_n + \tilde{x}_n)/2 \leq h_{n+1}.$$

5. THE KY FAN INEQUALITIES FOR A CERTAIN FAMILY OF MEANS

Let $X \in \mathbb{R}_+^n$. In [6] the authors have studied some unweighted means of X . They are denoted by A_n^+ , G_n^+ and H_n^+ and defined by $A_n^+ = A_n(X+1)$, $G_n^+ = G_n(X+1)$ and $H_n^+ = H_n(X+1)$, where $X+1 = (x_1+1, \dots, x_n+1)$. In the abovementioned paper, the authors have proven that

$$(5.1) \quad \frac{H_n}{H_n^+} \leq \frac{G_n}{G_n^+} \leq \frac{A_n}{A_n^+}.$$

Recently, Govedarica and Jovanović [9] have shown that

$$(5.2) \quad \frac{1}{H_n^+} - \frac{1}{H_n} \leq \frac{1}{G_n^+} - \frac{1}{G_n} \leq \frac{1}{A_n^+} - \frac{1}{A_n}.$$

It is worth mentioning that the inequalities (5.2) follow immediately from (5.1) and Lemma 2.1, part (i), because $A_n/A_n^+ \leq 1$.

Assume now that $x_i > 1$, $1 \leq i \leq n$ and let us define the unweighted means

$$A_n^- = A_n(X - 1), \quad G_n^- = G_n(X - 1), \quad H_n^- = H_n(X - 1),$$

where $X - 1 = (x_1 - 1, \dots, x_n - 1)$. The main result of this section reads as follows.

THEOREM 5.1. *The following inequalities*

$$(5.3) \quad 1 \leq \frac{A_n}{A_n^-} \leq \frac{G_n}{G_n^-} \leq \frac{H_n}{H_n^-}$$

and

$$(5.4) \quad \frac{1}{A_n^-} - \frac{1}{A_n} \leq \frac{1}{G_n^-} - \frac{1}{G_n} \leq \frac{1}{H_n^-} - \frac{1}{H_n}$$

hold true.

PROOF: The first inequality in (5.3) is an immediate consequence of the trivial identity $A_n = A_n^- + 1$. In order to prove the second inequality in (5.3) we use the following inequality ([10, Theorem 64], [12, 3.2.34, p. 208])

$$\prod_{i=1}^n (1 + y_i)^{1/n} \geq 1 + \prod_{i=1}^n y_i^{1/n}$$

with $y_i = x_i - 1$ ($1 \leq i \leq n$) to obtain $G_n \geq 1 + G_n^-$. This in conjunction with $G_n \leq A_n$ and $A_n^- = A_n - 1$ gives

$$\frac{G_n}{G_n^-} \geq \frac{G_n}{G_n - 1} \geq \frac{A_n}{A_n - 1} = \frac{A_n}{A_n^-}.$$

The last inequality in (5.3) is a special case of the following one

$$(5.5) \quad \frac{G_n(X)}{G_n(X - a)} \leq \frac{H_n(X)}{H_n(X - a)},$$

where now X is an n -tuple of positive numbers such that $\min(X) > a > 0$. Inequality (5.5) is reversed if $a < 0$. In order to establish (5.5) we define a function $f(x) = \prod_{i=1}^n x_i^{1/n}$ ($x_i > 0$ for $1 \leq i \leq n$). It is known that $f(x)$ is a concave function on \mathbb{R}_+^n . Thus its gradient ∇f satisfies the inequality

$$(\nabla f(X) - \nabla f(Y)) \cdot (X - Y) \leq 0$$

($X, Y \in \mathbb{R}_+^n$). Since $\nabla f(X) = (1/n) \prod_{i=1}^n x_i^{1/n} ((1/x_1), \dots, (1/x_n))$, the last inequality can be written as

$$\sum_{k=1}^n \left(\frac{G_n(X)}{x_k} - \frac{G_n(Y)}{y_k} \right) (x_k - y_k) \leq 0.$$

Letting $Y = X - a$ ($a > 0$) we obtain the desired result. \square

We close this section with a remark that the first inequality in (5.1) follows from (5.5) by letting $a = -1$.

6. INEQUALITIES INVOLVING MEANS OF TWO VARIABLES

This section deals with the Ky Fan inequalities for certain unweighted means of two variables $x > 0$ and $y > 0$. To this end we shall always assume that $x \neq y$. For the sake of notation we shall often omit symbols of the variables of a mean. Thus A will stand for the unweighted arithmetic mean of x and y , et cetera.

Recall that the identric I mean of order one of two variables is defined as

$$(6.1) \quad I \equiv I(x, y) = \exp\left(-1 + \frac{x \ln x - y \ln y}{x - y}\right).$$

The extended logarithmic mean E_t of order $t \in \mathbb{R}$ is defined as follows

$$(6.2) \quad E_t \equiv E_t(x, y) = \begin{cases} \left[\frac{x^t - y^t}{t(x - y)}\right]^{1/(t-1)} & t \neq 0, 1 \\ L(x, y), & t = 0 \\ I(x, y), & t = 1. \end{cases}$$

The latter mean is a special case of the two-parameter family of means $S_{p,q}$ ($p, q \in \mathbb{R}$) introduced by Stolarsky in [23]

$$S_{p,q} \equiv S_{p,q}(x, y) = \begin{cases} \left(\frac{q x^p - y^p}{p x^q - y^q}\right)^{1/(p-q)} & pq(p - q) \neq 0 \\ [L(x^p, y^p)]^{1/p}, & p \neq 0, q = 0 \\ [I(x^p, y^p)]^{1/p}, & p = q \neq 0 \\ G(x, y), & p = q = 0. \end{cases}$$

It is easy to see that

$$(6.3) \quad E_t = S_{t,1}.$$

Another two-parameter family of means, denoted by $G_{p,q}$, has been introduced by Gini in [8]. They are denoted by $G_{p,q}$ and defined as follows

$$G_{p,q} \equiv G_{p,q}(x, y) = \begin{cases} \left(\frac{x^p + y^p}{x^q + y^q}\right)^{1/(p-q)}, & p \neq q \\ [J(x^p, y^p)]^{1/p}, & p = q \neq 0 \\ G(x, y), & p = q = 0, \end{cases}$$

where

$$(6.4) \quad J(x, y) = \exp\left(\frac{x \ln x + y \ln y}{x + y}\right).$$

Other means used in this section are denoted by s_k and σ_k ($k = 1, 2, \dots$) and they are defined as follows

$$(6.5) \quad s_k \equiv s_k(x, y) = G^{1/k} E_{1/k}^{1-1/k}$$

and

$$(6.6) \quad \sigma_k \equiv \sigma_k(x, y) = G^{1/k} M_{1/k}^{1-1/k}.$$

Easy computations show that

$$(6.7) \quad s_k(x, y) = \frac{1}{k} \sum_{i=1}^k (x^{\alpha_i} y^{\beta_i} + x^{\beta_i} y^{\alpha_i}),$$

where $\alpha_i = (2i - 1)/2k$ and $\beta_i = 1 - \alpha_i$, $1 \leq i \leq k$. Also,

$$(6.8) \quad \sigma_1 = s_1 = G, \quad \sigma_2 = s_2 = (GM_{1/2})^{1/2} = \frac{1}{2}(x^{1/4}y^{3/4} + x^{3/4}y^{1/4}).$$

We shall also use the Heronian mean which will be denoted by He and is defined as

$$(6.9) \quad \text{He} \equiv \text{He}(x, y) = \frac{x + (xy)^{1/2} + y}{3}.$$

It is easy to verify that $\text{He} = S_{3/2, 1/2}$.

The following result will be used in the sequel.

PROPOSITION 6.1. *Let $\lambda = (t/2) \ln(x/y)$, $t \in \mathbb{R}$. Then*

$$(6.10) \quad \frac{\sinh \lambda}{\lambda} = \frac{LE_t^{t-1}}{G^t}$$

and

$$(6.11) \quad \cosh \lambda = \left(\frac{M_t}{G}\right)^t.$$

In particular, if $t = 1/k$ ($k = 1, 2, \dots$), then

$$(6.12) \quad \frac{\sinh \lambda}{\lambda} = \frac{L}{s_k}$$

and

$$(6.13) \quad \cosh \lambda = \frac{M_{1/k}}{\sigma_k}.$$

PROOF: Formulas (6.1) and (6.11) are valid when $t = 0$. Let $t \neq 0$. We have

$$\begin{aligned} \frac{\sinh \lambda}{\lambda} &= \frac{e^\lambda - e^{-\lambda}}{2\lambda} = \frac{(x/y)^{t/2} - (y/x)^{t/2}}{t \ln(x/y)} = \frac{(xy)^{t/2} [(x/y)^{t/2} - (y/x)^{t/2}]}{t \ln(x/y)(xy)^{t/2}} \\ &= \frac{x^t - y^t}{t \ln(x/y)(xy)^{t/2}} = \frac{x - y}{\ln(x/y)} \frac{x^t - y^t}{t(x - y)(xy)^{t/2}} = \frac{LE_t^{t-1}}{G^t}. \end{aligned}$$

Similarly,

$$\begin{aligned} \cosh \lambda &= \frac{1}{2}(e^\lambda + e^{-\lambda}) = \frac{1}{2}[(x/y)^{t/2} + (y/x)^{t/2}] \\ &= \frac{(xy)^{t/2} [(x/y)^{t/2} + (y/x)^{t/2}]}{2(xy)^{t/2}} = \frac{(x^t + y^t)/2}{(xy)^{t/2}} = \left(\frac{M_t}{G}\right)^t. \end{aligned}$$

For the proof of (6.12) we let $t = 1/k$ in (6.2) to obtain

$$E_{1/k}^{1/k-1} = \frac{k}{\sum_{i=1}^k x^{(k-i)/k} y^{(i-1)/k}}.$$

Making use of (6.7) we obtain

$$\frac{E_{1/k}^{1/k-1}}{G^{1/k}} = \frac{1}{s_k}.$$

This in conjunction with (6.10) gives the assertion (6.12). Formula (6.13) follows from (6.11) by letting $t = 1/k$ ($k \in \mathbb{N}$). We have

$$\begin{aligned} \cosh \lambda &= \frac{(x^{1/k} + y^{1/k})/2}{(xy)^{1/2k}} = \frac{(x^{1/k} + y^{1/k})(x^{1/k} + y^{1/k})^{k-1}/2^k}{(xy)^{1/2k}(x^{1/k} + y^{1/k})^{k-1}/2^{k-1}} \\ &= \frac{[(x^{1/k} + y^{1/k})/2]^k}{(xy)^{1/2k} [(x^{1/k} + y^{1/k})/2]^{k-1}} = \frac{M_{1/k}}{G^{1/k} M_{1/k}^{1-1/k}} = \frac{M_{1/k}}{\sigma_k}, \end{aligned}$$

where in the last step we have used formula (6.6). The proof is complete. \square

To this end we shall always assume that $0 < x, y \leq 1/2$. We are in a position to prove the following.

THEOREM 6.2. *Let $k = 1, 2, \dots$. Then the following inequalities*

$$(6.14) \quad \frac{H}{H'} < \frac{\sqrt{HG}}{\sqrt{H'G'}} < \frac{\sqrt{HL}}{\sqrt{H'L'}} < \frac{G}{G'} < \frac{s_2}{s'_2},$$

$$(6.15) \quad \frac{s_k}{s'_k} < \frac{L}{L'} < \frac{M_{1/2}}{M'_{1/2}} < \frac{\text{He}}{\text{He}'} < \frac{M_{2/3}}{M'_{2/3}} < \frac{I}{I'} < \frac{A}{A'} < \frac{M_2}{M'_2} < \frac{J}{J'},$$

and

$$(6.16) \quad \frac{G}{G'} \leq \frac{\sigma_k}{\sigma'_k} < \frac{M_{1/k}}{M'_{1/k}}$$

are valid with equality in (6.16) when $k = 1$.

PROOF: The first inequality in (6.14) follows from the Wang–Wang inequality (1.2) and Lemma 2.1, part (iii) while the second inequality in (6.14) follows from $G/G' < L/L'$ (see [7, 16]). Multiplying both sides by H/H' and next extracting the square roots, we obtain the desired result. For the proof of the third inequality in (6.14) we apply the Ky Fan inequality for the Gini and Stolarsky means

$$(6.17) \quad \frac{G_{p,q}}{G'_{p,q}} < \frac{S_{p,q}}{S'_{p,q}}$$

($p + q < 0$) (see [16]). Letting $p = -1$, $q = 0$ and next utilising definitions of the Gini means and Stolarsky means we obtain $G_{-1,0} = H$ and $S_{-1,0} = G^2/L$. Substituting into

(6.17) gives the assertion. The last inequality in (6.14) can be established as follows. We use $G/G' < M_{1/2}/M'_{1/2}$ (see, for example, [5, 16]). The first inequality in (2.4) together with $s_2 = (GM_{1/2})^{1/2}$ implies

$$\frac{G}{G'} < \left(\frac{GM_{1/2}}{G'M'_{1/2}} \right)^{1/2} = \frac{s_2}{s'_2}.$$

Here we have used the identity $M_{1/2} = E_{1/2}$. We shall establish now the chain of the inequalities (6.15). The first one follows by application of Lemma 2.3, with $p = (1/2k) \ln(x/y)$ and $q = (1/2k) \ln(x'/y')$ to (6.12). For the proof of the second inequality in (6.15) we combine (6.12) with (6.13) to obtain

$$\frac{\tanh \lambda}{\lambda} = \frac{L \sigma_k}{M_{1/k} s_k}.$$

Using (2.7) with p and q as defined earlier, we obtain

$$\frac{L \sigma_k}{L' \sigma'_k} < \frac{M_{1/k} s_k}{M'_{1/k} s'_k} \quad (k = 1, 2, \dots).$$

Letting above $k = 2$ and taking into account that $\sigma_2 = s_2$ (see (6.8)), we obtain the desired result. The third inequality in (6.15) can be proven as follows. Assume without a loss of generality that $x < y$ and next define $t^2 = y/x$ and $t'^2 = (1-y)/(1-x)$. Clearly $t > t'$. Also, let

$$f(t) = \frac{M_{1/2}(t^2, 1)}{\text{He}(t^2, 1)}.$$

An easy computation shows that $f(t) = \frac{3}{4} \left(1 + \frac{t}{t^2 + t + 1} \right)$ and also that $f'(t) = \frac{3}{4} \frac{1 - t^2}{(t^2 + t + 1)^2}$. Thus the function $f(t)$ is strictly decreasing when $t^2 > 1$. This in turn implies that

$$\frac{M_{1/2}(t^2, 1)}{\text{He}(t^2, 1)} < \frac{M_{1/2}(t'^2, 1)}{\text{He}(t'^2, 1)}.$$

Replacing t^2 and t'^2 by their defining expressions, next multiplying the numerators on both sides by x and the denominators by $1 - x$ and utilising homogeneity and symmetry of both means in their variables, we obtain the desired result. For the proof of the fourth inequality in (6.15) we still assume that $x < y$ and we let $t^3 = y/x$ and $t'^3 = (1-y)/(1-x)$. Again we have $t > t'$. Also, let

$$g(t) = \frac{\text{He}(t^3, 1)}{M_{2/3}(t^3, 1)}.$$

Logarithmic differentiation gives

$$\frac{g'(t)}{g(t)} = -\frac{3 t^{1/2}(t-1)(t^{1/2}-1)^2}{2(t^3+t^{3/2}+1)(t^2+1)} < 0.$$

Thus the positive function $g(t)$ is decreasing for $t > 1$. We follow the lines introduced in the proof of the previous inequality to obtain the assertion. The fifth inequality in (6.15) is established in a similar manner. With t and t' as in the proof of the last inequality and with

$$k(t) = \left[\frac{M_{2/3}(t^3, 1)}{I(t^3, 1)} \right]^{2/3}$$

we have

$$\frac{k'(t)}{k(t)} = \frac{2t^2}{(t^3 - 1)^2} s(t),$$

where

$$s(t) = 3 \ln t - \frac{(t+1)(t^3-1)}{t(t^2+1)}.$$

Letting $u = t^3$ in Karamata's inequality [12, 3.616, p. 272]

$$\frac{\ln u}{u-1} \leq \frac{1+u^{1/3}}{u+u^{1/3}}$$

we obtain $s(t) \leq 0$ for $t > 1$. Thus the function $k(t)$ is strictly decreasing. The fifth inequality in (6.15) now follows by using the same argument as in the proof of the last two previous inequalities. The sixth and seventh inequalities in (6.15) are known (see (1.12) and [5], respectively). To prove the last inequality in (6.15) we proceed as follows. Again assume that $x < y$ and put $t = y/x$ and $t' = (1-y)/(1-x)$. Also, let

$$f(t) = \frac{M_2(t, 1)}{J(t, 1)},$$

where the Gini mean J is defined in (6.4). We have

$$f(t) = \left(\frac{t^2+1}{2} \right)^{1/2} \frac{1}{t^{t/(t+1)}}.$$

Hence

$$\frac{f'(t)}{f(t)} = \frac{h(t)}{(t+1)^2(t^2+1)},$$

where $h(t) = t^2 - 1 - (t^2 + 1) \ln t$. In order to prove that the function $f(t)$ is strictly decreasing for $t > 1$ we use

$$(6.18) \quad \frac{t-1}{\ln t} = L(t, 1) < \frac{t+1}{2} < \frac{t^2+1}{t+1},$$

where the second inequality follows from the following one $(t+1)^2 < 2(t^2+1)$. It follows from (6.18) that $h(t) < 0$ for $t > 1$. This in turn implies that $f(t)$ is strictly decreasing on the stated domain. We shall prove now the inequalities (6.16). It follows from (6.6) that

$$\frac{\sigma_k}{G} = \left(\frac{M_{1/k}}{G} \right)^{1-1/k}.$$

On the other hand, letting $t = 1/k$ in (6.11) we obtain

$$\frac{M_{1/k}}{G} = (\cosh \lambda)^k.$$

Combining the last two formulas we obtain

$$\frac{\sigma_k}{G} = (\cosh \lambda)^{k-1}.$$

Next we use inequality (2.5) of Lemma 2.3 to obtain the desired result. The second inequality in (6.16) is also proven with the aid of Lemma 2.3 and formula (6.13). We omit further details. The proof is complete. \square

More Ky Fan type inequalities can be obtained immediately from (6.14)–(6.16) by using the Ky Fan Rules (i)–(iii) of Lemma 2.1. We omit further details.

Our next result reads as follows.

THEOREM 6.3. *The following inequalities*

$$(6.19) \quad \left(\frac{M_{1/2} G^3}{M'_{1/2} G'^3} \right)^{1/2} < \frac{L \operatorname{He}}{L' \operatorname{He}'} < \left(\frac{M_{1/2}}{M'_{1/2}} \right)^{1/2} \left(\frac{M_{3/2}}{M'_{3/2}} \right)^{3/2}$$

hold true.

PROOF: We use (6.10) with $t = 3/2$ and also utilise $E_{3/2}^{1/2} = \operatorname{He} / M_{1/2}^{1/2}$ (see (6.2)) to obtain

$$\frac{\sinh \lambda}{\lambda} = \frac{L \operatorname{He}}{(M_{1/2} G^3)^{1/2}},$$

where now $\lambda = (3/4) \ln(x/y)$. Application of the inequality (2.5) with $p = (3/4) \ln(x/y)$ and $q = (3/4) \ln(x'/y')$, completes the proof of the first inequality in (6.19). In order to establish the second one we combine (6.10) with (6.11) to obtain

$$\frac{\tanh \lambda}{\lambda} = \frac{L E_t^{t-1}}{M_t^t},$$

where now $\lambda = (t/2) \ln(x/y)$. Making use of Lemma 2.3 we obtain

$$\frac{L E_t^{t-1}}{M_t^t} < \frac{L'(E'_t)^{t-1}}{(M'_t)^t}.$$

Hence

$$\frac{L}{L'} \left(\frac{E_t}{E'_t} \right)^{t-1} < \left(\frac{M_t}{M'_t} \right)^t.$$

Letting $t = 3/2$ and using $E_{3/2}^{1/2} = \operatorname{He} / M_{1/2}^{1/2}$ we obtain the assertion. \square

Neuman and Sándor [15] have proven that

$$\frac{1}{H'} - \frac{1}{L'} < \frac{1}{H} - \frac{1}{L}.$$

The following result provides an improvement of the last inequality. We have

THEOREM 6.4. *The following inequality*

$$(6.20) \quad A'\left(\frac{1}{H'} - \frac{1}{L'}\right) < A\left(\frac{1}{H} - \frac{1}{L}\right)$$

is valid.

PROOF: We shall use the following series expansion

$$(6.21) \quad A\left(\frac{1}{H} - \frac{1}{L}\right) = \sum_{k=1}^{\infty} \frac{2k}{2k+1} \left(\frac{x-y}{x+y}\right)^{2k}.$$

For its proof we employ (2.9) with $a = x$, $b = y$, and $f(t) = 1/t$. Utilising the formula

$$\frac{1}{L} = \frac{1}{x-y} \int_y^x \frac{dt}{t}$$

we obtain

$$\frac{1}{L} = \frac{1}{H} - \frac{1}{A} \sum_{k=1}^{\infty} \frac{2k}{2k+1} \left(\frac{x-y}{x+y}\right)^{2k}.$$

Hence the formula (6.21) follows. For the proof of (6.20) we let

$$u = \frac{x-y}{x+y} \quad \text{and} \quad v = \frac{x'-y'}{x'-y'},$$

where $x' = 1-x$ and $y' = 1-y$. One can easily verify that $|v| \leq |u|$. Making use of (6.21) we obtain

$$A'\left(\frac{1}{H'} - \frac{1}{L'}\right) = \sum_{k=1}^{\infty} \frac{2k}{2k+1} v^{2k} \leq \sum_{k=1}^{\infty} \frac{2k}{2k+1} u^{2k} = A\left(\frac{1}{H} - \frac{1}{L}\right).$$

The proof is complete. □

REFERENCES

- M [1] H. Alzer, 'Inequalities for arithmetic, geometric and harmonic means', *Bull. London Math. Soc.* **22** (1990), 362–366.
- M [2] H. Alzer, 'The inequality of Ky Fan and related results', *Acta Appl. Math.* **38** (1995), 305–354.
- M [3] E.F. Beckenbach and R. Bellman, *Inequalities* (Springer-Verlag, New York, 1965).
- M [4] B.C. Carlson, *Special functions of applied mathematics* (Academic Press, New York, 1977).
- M [5] F. Chan, D. Goldberg and S. Gonek, 'On extensions of an inequality among means', *Proc. Amer. Math. Soc.* **42** (1974), 202–207.
- M [6] E. El-Newehi and F. Proschan, 'Unified treatment of some inequalities among means', *Proc. Amer. Math. Soc.* **81** (1981), 388–390.
- M [7] I. Gavrea and T. Trif, 'On Ky Fan's inequality', *Math. Inequal. Appl.* **4** (2001), 223–230.

- ^ [8] G. Gini, ‘Di una formula comprensiva delle medie’, *Metron* **13** (1938), 3–22.
- ^ [9] V. Govedarica and M. Jovanović, ‘On the inequalities of Ky Fan, Wang–Wang and Alzer’, *J. Math. Anal. Appl.* **270** (2002), 709–712.
- ℳ [10] G.H. Hardy, J.E. Littlewood and G. Pólya, *Inequalities*, (2nd edition) (Cambridge Univ. Press, London and New York, 1952).
- ℳ [11] N. Levinson, ‘Generalization of an inequality of Ky Fan’, *J. Math. Anal. Appl.* **8** (1964), 133–134.
- ^ [12] D.S. Mitrinović, *Analytic inequalities* (Springer-Verlag, Berlin, 1970).
- ℳ [13] E. Neuman, ‘Bounds for symmetric elliptic integrals’, *J. Approx. Theory* **122** (2003), 249–259.
- ℳ [14] E. Neuman, C.E.M. Pearce, J. Pečarić and V. Šimić, ‘The generalized Hadamard inequality, g -convexity and functional Stolarsky means’, *Bull. Austral. Math. Soc.* **68** (2003), 303–316.
- ℳ [15] E. Neuman and J. Sándor, ‘On the Ky Fan inequality and related inequalities I’, *Math. Inequal. Appl.* **5** (2002), 49–56.
- ℳ [16] E. Neuman and J. Sándor, ‘Inequalities involving Stolarsky and Gini means’, *Math. Pannon.* **14** (2003), 29–44.
- ℳ [17] E. Neuman and J. Sándor, ‘On the Schwab–Borchardt mean’, *Math. Pannon.* **14** (2003), 253–266.
- ℳ [18] J. Pečarić and V. Šimić, ‘Stolarsky–Tobey mean in n variables’, *Math. Inequal. Appl.* **2** (1999), 325–341.
- ^ [19] J. Sándor, ‘Sur la fonction Gamma’, *Publ. Centre. Rech. Math. Pures., Neuchâtel, Ser. I* **21** (1989), 4–7.
- ℳ [20] J. Sándor, ‘On an inequality of Ky Fan’, (Preprint 90-7, Babes-Bolyai Univ.), pp. 29–34.
- ℳ [21] J. Sándor, ‘On the identric and logarithmic means’, *Aequationes Math.* **40** (1990), 261–270.
- ℳ [22] J. Sándor and T. Trif, ‘A new refinement of the Ky Fan inequality’, *Math. Inequal. Appl.* **2** (1999), 529–533.
- ^ [23] K.B. Stolarsky, ‘Generalizations of the logarithmic mean’, *Math. Mag.* **48** (1975), 87–92.
- ℳ [24] W.-L. Wang and P.-F. Wang, ‘A class of inequalities for symmetric functions’, (in Chinese), *Acta Math. Sinica* **27** (1984), 485–497.

Department of Mathematics
 Southern Illinois University
 Carbondale, IL 62901-4408
 United States of America
 e-mail: edneuman@math.siu.edu
 urladdr:
<http://www.math.siu.edu/neuman/personal.html>

Department of Pure Mathematics
 Babes-Bolyai University
 Ro-3400 Cluj-Napoca
 Romania
 e-mail: jsandor@math.ubbcluj.ro