

## A correspondence note on Myerson's 'Irrationality via well-ordering'

Scott Duke Kominers\*

I thoroughly enjoyed Myerson's article [2] on methods of proving irrationality via the well-ordering principle. In this note, I point out a second method by which the well-ordering principle may be used to prove Myerson's Theorem 2. This approach is a generalisation of MacHale's recent proof [1] that  $\sqrt{2}$  is irrational.

**Theorem 1.** *For  $m \in \mathbb{N}$ ,  $\sqrt{m}$  is irrational if it is not an integer.*

*Proof.* First, we prove the result for squarefree  $m$ . We consider the set

$$\{a + b \mid a, b \in \mathbb{N}, a^2 = mb^2\}.$$

By the well-ordering principle for  $\mathbb{N}$ , this set has a minimal element  $a_0 + b_0$ . If  $m$  is squarefree, the condition  $a_0^2 = mb_0^2$  guarantees that  $m$  divides  $a_0$ . Thus,  $a_0 = m\ell$  for some  $\ell \in \mathbb{N}$ . But then,  $m^2\ell^2 = a_0^2 = mb_0^2$ , from which it follows that  $m$  divides  $b_0$ . Writing  $b_0 = mr$ , we have  $m^3r^2 = m^2\ell^2$ . It follows that  $mr^2 = \ell^2$ . As  $a_0 + b_0 = m(\ell + r)$ , this is a contradiction to the minimality of  $a_0 + b_0$ .

Now, if  $m$  is not squarefree, we may write  $m = m_1^2 m_0$  for  $m_0$  squarefree. We have  $\sqrt{m_0}$  irrational by the earlier argument, so  $\sqrt{m} = \sqrt{m_1^2 m_0} = m_1 \sqrt{m_0}$  is irrational, as well.

### References

- [1] MacHale, D. (2008). The well-ordering principle for  $\mathbb{N}$ . *Math. Gaz.* **92**, 257–259.  
 [2] Myerson, G. (2008). Irrationality via well-ordering. *Gaz. Aust. Math. Soc.* **35**, 121–125.

---

Received 18 July 2008; accepted for publication 3 December 2008.

\*Student, Department of Mathematics, Harvard University

c/o 8520 Burning Tree Road, Bethesda, MD 20817, USA. E-mail: kominers@fas.harvard.edu